

Plausible Sources of Membrane-Forming Fatty Acids on the Early Earth: A Review of the Literature and an Estimation of Amounts

Zachary R. Cohen,* Zoe R. Todd, Nicholas Wogan, Roy A. Black, Sarah L. Keller, and David C. Catling*



Cite This: *ACS Earth Space Chem.* 2023, 7, 11–27




Read Online

ACCESS |

Metrics & More

Article Recommendations

ABSTRACT: The first cells were plausibly bounded by membranes assembled from fatty acids with at least 8 carbons. Although the presence of fatty acids on the early Earth is widely assumed within the astrobiology community, there is no consensus regarding their origin and abundance. In this Review, we highlight three possible sources of fatty acids: (1) delivery by carbonaceous meteorites, (2) synthesis on metals delivered by impactors, and (3) electrochemical synthesis by spark discharges. We also discuss fatty acid synthesis by UV or particle irradiation, gas-phase ion–molecule reactions, and aqueous redox reactions. We compare estimates for the total mass of fatty acids supplied to Earth by each source during the Hadean eon after an extremely massive asteroid impact that would have reset Earth's fatty acid inventory. We find that synthesis on iron-rich surfaces derived from the massive impactor in contact with an impact-generated reducing atmosphere could have contributed $\sim 10^2$ times more total mass of fatty acids than subsequent delivery by either carbonaceous meteorites or electrochemical synthesis. Additionally, we estimate that a single carbonaceous meteorite would not deliver a high enough concentration of fatty acids (~ 15 mM for decanoic acid) into an existing body of water on the Earth's surface to spontaneously form membranes unless the fatty acids were further concentrated by another mechanism, such as subsequent evaporation of the water. Our estimates rely heavily on various assumptions, leading to significant uncertainties; nevertheless, these estimates provide rough order-of-magnitude comparisons of various sources of fatty acids on the early Earth. We also suggest specific experiments to improve future estimates. Our calculations support the view that fatty acids would have been available on the early Earth. Further investigation is needed to assess the mechanisms by which fatty acids could have been concentrated sufficiently to assemble into membranes during the origin of life.



KEYWORDS: Fatty acid, membrane, prebiotic chemistry, origin of life, astrobiology

INTRODUCTION

Cells use bilayer membranes to separate themselves from their environment. In modern cells, these membranes are composed of phospholipids. During the origin of cells on Earth, more primitive membranes likely played a similar role,¹ sequestering cellular building blocks^{2,3} and polymers.⁴ The hydrocarbons in modern phospholipids are tails of fatty acids connected by an ester linkage to the glycerol backbone. Fatty acids themselves can assemble into membranes. Fatty acids were likely more abundant than phospholipids on the early Earth, leading to a common hypothesis that the membranes of the first cells were composed of fatty acids.¹

Fatty acids consist of a hydrocarbon tail and a carboxylic acid headgroup (Figure 1). Saturated fatty acids with eight or more carbons in a linear chain can assemble into membranes.⁵ Fatty acids with unsaturated⁶ or branched⁷ chains can also assemble into membranes, although in these cases it is unknown whether the minimum number of carbons for

membrane assembly is greater or less than eight. Fatty acids with carboxylic acids at both ends of a carbon chain do not assemble into membranes on their own, although these dicarboxylic acids can incorporate into membranes when additional types of amphiphiles are present.⁸


To form membranes, the solution pH must be within about half a unit of the effective pK_a of the fatty acid in a bilayer⁵ (where $pK_a = -\log_{10}$ of the equilibrium constant, K_a , for the dissociation of the fatty acid into a proton and the negatively charged amphiphile). If the solution pH is high enough such that the vast majority of headgroups are charged or if the fatty

Received: June 1, 2022

Revised: October 18, 2022

Accepted: October 24, 2022

Published: December 22, 2022






Figure 1. Fatty acid assembly depends on the pH of the surrounding solution. When the pH is below the effective pK_a of the fatty acids in a bilayer, fatty acids form an oil that is immiscible with the surrounding aqueous solution (bottom). When the pH is near the effective pK_a , fatty acids assemble into bilayers in a membrane (middle). Vesicles, spherical shells of these membranes, may have served as the membrane compartments for the first cells on Earth. When the pH is above the pK_a of the fatty acids in a bilayer, the fatty acids assemble into micelles, which cannot encapsulate aqueous solutes (top).

acids have fewer than eight carbons, then fatty acids assemble into nanoscale micelles that cannot encapsulate aqueous solutes (Figure 1). On the other hand, if the solution pH is low enough that the vast majority of headgroups are uncharged, then fatty acids separate into an oil phase.⁹

Fatty acid membranes provide some advantages for early cell replication compared to modern phospholipid membranes.⁹ For example, fatty acid membranes are moderately permeable to salts and small organic molecules such as nucleotides, allowing internal replication of nucleic acids, which would have been critical for developing cells.¹⁰ The surface area of these vesicles increases when they incorporate additional fatty acids from the environment into the membrane.^{11,12} A growing vesicle can be supplied with fatty acids from micelles^{11,13} or from other vesicles.¹⁴ Vesicles that retain fatty acids grow while others shrink, which could have enabled competition between primitive cells for a limited supply of fatty acids.¹² After acquiring excess membrane surface area, primitive cells could have divided when exposed to modest shear stress.^{13,15} Vesicles of phospholipids do not grow or divide as readily because aqueous solubility of a phospholipid with two hydrophobic tails is much lower than the solubility of a single-tailed fatty acid, so transfer through aqueous solution is slower.^{16,17}

Here, we review how membrane-forming fatty acids could have been supplied on the early Earth. We identify three relatively well-characterized sources of abiotic fatty acids: delivery by meteorites, synthesis on the surface of metal catalysts, and synthesis by electrochemistry (Figure 2). There are also reports of fatty acid syntheses that fall outside these categories, but these have been less robustly investigated. We summarize the important details of the experiments that have been carried out and discuss whether similar reactions could have plausibly occurred on the early Earth. Finally, we estimate




Figure 2. There are three well-characterized sources that could have provided fatty acids to the early Earth. (A) Carbonaceous meteorites can deliver fatty acids.^{19–22} (B) Metal surfaces can catalyze fatty acid synthesis. As one example, Noonan and Oro mixed filings of the Canyon Diablo meteorite (containing iron and nickel) with deuterium and carbon monoxide gases, and the mixture was heated to 400 °C to produce fatty acids.²³ Similar experiments have used pure Fe, Ni, or Fe- and Ni-containing minerals as catalysts and a variety of carbon and hydrogen sources to synthesize fatty acids (Table 1). (C) Fatty acids can also be synthesized during electrical sparging (Table 2). As one example, Yuen et al. used an electric discharge to synthesize fatty acids from methane.²⁴

the total mass of fatty acids produced from each of the three relatively well-characterized sources during the Hadean eon when the first cells are hypothesized to have formed.¹⁸

■ CARBONACEOUS METEORITES DELIVER FATTY ACIDS TO EARTH

Fatty acids have been detected in a variety of carbonaceous meteorites that have landed on Earth,^{19–22} and molecules extracted from at least one such meteorite assemble into membranes.²⁵ At least 18 different carbonaceous meteorites have been analyzed, and both linear-chain and branched-chain fatty acids have been identified.²⁶ Depending on the type of carbonaceous meteorite, the abundance of membrane-forming fatty acids can range from 1 ppb to 100 ppm by mass.²⁶ Recent reviews provide detailed analyses of meteoritic fatty acids.^{26,27} Importantly, it remains unclear which types of reactions are responsible for synthesizing meteoritic fatty acids in space.²⁶

Could fatty acids delivered by carbonaceous meteorites have dissolved into water and assembled into membranes? Meteorites can fragment in airbursts during passage through the atmosphere, allowing some fatty acids to remain intact.^{28,29} Meteorites with radii less than 100 m tend to fragment when differential pressure across the small body exceeds the material strength.²⁸ The Chelyabinsk ordinary chondrite meteorite (radius ~ 10 m) that fell in Russia in 2013 fragmented at an altitude above 25 km, and fragments were spread into an area 250–300 km² around the trajectory.³⁰ The largest fragment

was ~ 0.7 m mean diameter and fell into a lake. Fatty acids in carbonaceous chondrites would likely be similarly dispersed over a wide spatial area.

When meteorite fragments disperse into water on Earth's surface, fatty acids can dissolve. Numerical models suggest that nucleobases leach out of 20 cm meteorite fragments and mix into the surrounding water within about three years.²⁹ At moderately alkaline pH, fatty acids can be even more soluble because of their charged headgroup. However, in order for fatty acids to assemble into membranes, the fatty acids must accumulate in solution above the critical vesicle concentration (~ 15 mM for decanoic acid³). If fatty acids are present at concentrations below the critical vesicle concentration, membrane assembly does not occur.

Here, we use the measured abundance of decanoic acid (a 10-carbon fatty acid) in carbonaceous meteorites²⁶ to calculate the volume of a meteorite fragment that would be required to deliver enough decanoic acid into water so that the concentration of decanoic acid equals the critical vesicle concentration. Although it is not known precisely how meteorite size, initial velocity, or impact angle influences the fraction of fatty acids that survive impact, we note that a large portion of the meteorite's initial mass (and thus a large portion of its fatty acids) may be destroyed by ablation and heating during travel through the atmosphere.^{31–33} Given that our estimates rely on the average mass fraction of decanoic acid that has been recovered from natural carbonaceous meteorites and that we do not know a priori the mass, velocity, and impact angle of each meteorite, we assume the measured fatty acid abundances represent the average survival over the entire population of possible impacts. Although this assumption introduces potential errors, a more precise calculation is beyond the scope of this study.

The volume of a meteorite fragment (V_{meteor} , expressed in m^3) that would be required to deliver enough decanoic acid to reach the critical vesicle concentration as a function of water volume (V_{water} , expressed in m^3) is given by eq 1:

$$V_{\text{meteor}} = \frac{C_{\text{cvc}} V_{\text{water}} w}{a \rho} \approx 61.5 V_{\text{water}} \quad (1)$$

where C_{cvc} is the critical vesicle concentration for decanoic acid (15 mol/m^3 of water, equivalent to 15 mM), w is the molar mass of decanoic acid (0.1723 kg/mol), a is the dimensionless fraction of the meteorite's mass that is decanoic acid (2×10^{-5} for CM2 type meteorites), and ρ is the meteorite density (2100 kg/m^3 for CM type meteorites).

The result of our estimate is shown in Figure 3. To deliver enough decanoic acid to form membranes, the volume of the meteorite fragment would have to exceed the volume of the waterbody. If a meteorite fragment were to land in a large enough waterbody to submerge the fragment, the decanoic acid that subsequently dissolved in the water would be too dilute to form membranes. However, this does not rule out membrane formation. The concentration of fatty acids could increase during dry periods, accompanied by a net loss of water due to evaporation.³⁴ Although we limited our calculation to decanoic acid because the critical vesicle concentration has been measured, a range of fatty acids from 8 to 12 carbons can be delivered simultaneously during a meteorite impact. The presence of these additional fatty acids could enable membrane formation at lower concentrations of decanoic acid.^{35,36} We do not consider that CM type meteorites can contain significant water (up to $\sim 9\%$ by mass³⁷). If all meteoritic fatty acids were




Figure 3. A single fragment of a carbonaceous meteorite cannot directly deliver enough decanoic acid to a body of water to form membranes. To exceed the critical vesicle concentration ($\sim 15 \text{ mM}$),³ the volume of the meteorite (red line) would exceed the volume of the water. However, subsequent evaporation of the water could concentrate decanoic acid and enable membrane formation. A CM2 type meteorite is assumed because it contains the most decanoic acid on average (20 ppm by mass²⁶). The density of CM-type meteorites is 2100 kg/m^3 .³⁸ Only meteorites with radii less than 100 m ($\sim 10^6 \text{ m}^3$ for a spherical meteorite) can fragment and impact the Earth's surface with low enough energy to preserve fatty acids.^{28,29} See eq 1 for details of the calculation.

to somehow dissolve in only this meteoritic water, the concentration of decanoic acid could exceed the critical vesicle concentration; this can be interpreted as an upper limit for the concentration of fatty acids delivered by a meteorite.

To conclude our section on carbonaceous meteorites, we find that meteorite delivery was unlikely to directly yield high enough aqueous concentrations of fatty acids to form membranes on the early Earth. Additional processes would have been necessary to further concentrate fatty acids above the critical vesicle concentration, which we cannot rule out.

■ CATALYSIS OF FATTY ACID SYNTHESIS BY METAL SURFACES

The most commonly reported abiotic synthesis of fatty acids involves catalytic metal surfaces. Within this class of syntheses, Fischer–Tropsch reactions are the most thoroughly investigated. Fischer–Tropsch reactions occur when H_2 and either CO or CO_2 adsorb onto a metal surface.^{39,40} Surfaces of solid iron or nickel are most commonly tested, although FeS , NiS , and Fe_3O_4 minerals have also been used to produce fatty acids.^{41–44} In most experiments, metal surfaces must be heated above $150 \text{ }^\circ\text{C}$ to produce fatty acids. Catalysts contain metals in their reduced form; the synthesis of membrane-forming fatty acids (at least 8 carbons) has not been demonstrated on oxidized metal surfaces.^{23,45–48} The synthesis of fatty acids

seems to occur at the gas–solid interface (Figure 4). Even in experiments designed to eliminate gaseous headspace,




Figure 4. Fatty acid synthesis occurs at the interfaces between reducing metal surfaces and a gaseous headspace. (A) Nooner and Oro showed that deuterium and carbon monoxide gases react together on the surface of hot (400 °C) meteorite filings to produce membrane-forming fatty acids.^{23,50} When the meteorite filings were artificially oxidized, fatty acid synthesis was not observed.²³ (B) In hydrothermal experiments, McCollum et al. report that the synthesis of membrane-forming fatty acids occurs within gaseous bubbles adsorbed onto oxidation-resistant stainless steel surfaces.⁴⁹ When oxidized metal surfaces are present instead of stainless steel, only short-chain (<5 carbons) fatty acids are formed.^{23,45–48} In these hydrothermal experiments, aqueous formic acid or oxalic acid is used for experimental convenience as a source of H₂ and CO₂.

reactions are suggested to proceed in gaseous bubbles within the aqueous solution.⁴⁸ Whether or not an explicit gaseous headspace is present, the carbon-containing precursors are generally supplied as gases. However, for experimental convenience in hydrothermal experiments, both formic acid and oxalic acid have been used as aqueous starting materials for fatty acid synthesis because both compounds decompose into H₂ and CO₂ at temperatures above 150 °C.⁴⁹

Metal-catalyzed reactions generally create a diverse mixture of product types, including hydrocarbons and fatty alcohols in addition to fatty acids.^{48,49,51–54} For each type of product, molecules with more carbons are less abundant.⁴⁰ Many experiments have produced only short-chain carboxylic acids

containing fewer than the 8 carbons required for membrane assembly⁵ (Figure 5). The carbon chain of a fatty acid could be elongated upon further reaction with a metal catalyst,⁴⁰ although additional experiments are required to validate this hypothesis. Table 1 summarizes experiments in the literature that have produced fatty acids using metal catalysts.

Metal-catalyzed reactions could have occurred on the early Earth after meteorite impacts, which delivered iron, nickel, and heat. Meteoritic metals could have been exposed to atmospheric H₂, CO₂, and CO, enabling surface-catalyzed synthesis of fatty acids. Extremely large impactors, about the size of the asteroid Vesta (~10²⁰ kg), could have transformed the Earth into a global Fischer–Tropsch reactor with surface temperatures >100 °C and high partial pressures of H₂, CO₂, and CO for thousands of years.⁵⁵ Catalytic metal surfaces would be required to produce fatty acids, so future research in this area will be especially valuable if it constrains the location and quantity of reduced metals after such impacts.⁵⁶ Although a large impact would have been catastrophic for any life that was already present on Earth, it could have potentially seeded Earth's postimpact surface with fatty acids and other necessary biomolecules, enabling life to subsequently emerge.⁵⁷

Ground-breaking experiments by Nooner and Oro modeled a postimpact scenario for fatty acid synthesis.^{23,50} It remains uncertain how the yield of fatty acids depends on experimental parameters such as partial pressure and temperature. Kinetic models for the Fischer–Tropsch process have been developed in industrial settings, which generally do not mimic plausible early Earth conditions, and the quantitative form of the models depends on the design of the reactor.⁵⁸ Until experiments are conducted to understand how fatty acid production depends on reaction parameters more relevant to the early Earth, there will be substantial uncertainty in estimates of production of fatty acids by metal catalysts on the early Earth.

Another natural setting in which the metal catalyzed synthesis of fatty acids might occur is in a hydrothermal environment, where the conversion of CO₂ and H₂ into fatty acids is thermodynamically favorable.⁵⁹ Fatty acids have been detected in natural hydrothermal systems; however, it is unclear what fraction of those fatty acids were produced by




Figure 5. Length of fatty acids delivered by meteorites (labeled “Meteor.”) or produced in abiotic synthesis experiments. Fatty acids with at least eight carbons, indicated by the dashed red line, can assemble into membranes. All the fatty acids produced are saturated and unbranched (except in Scheidler et al. 2016, where experiments also produced unsaturated fatty acids). Note that detection of a fatty acid with a certain length may not have been attempted during every experiment.

Table 1. Summary of Experiments That Used Metal Surfaces to Catalyze the Synthesis of Fatty Acids^a

year and ref	explicit gas phase?	explicit aqueous phase?	H ₂ source	carbon source	solid surface	reaction conditions	number of carbons per fatty acid
1976 ⁵⁰	Yes	No	D ₂ (g)	CO (g)	Meteorite filings and K ₂ CO ₃	50 h at 370 °C	10–16
1979 ²³	Yes	No	D ₂ (g)	CO (g)	Meteorite filings (containing iron and nickel) and carbonate salts	48 h at 400 °C	5–21
1997 ⁴¹	Yes	Yes	N/A	CO (g) and CH ₃ SH (aq)	NiS–FeS	1 week at 100 °C	2
1999 ⁴⁹	Yes	Yes	Formic acid or oxalic acid (aq)	Formic acid or oxalic acid (aq)	Stainless steel	>48 h at 175 °C	6–22
2001 ⁵¹	No	Yes	Oxalic acid (aq)	Oxalic acid (aq)	Stainless steel	48 h at 100 °C	7–18
2003 ⁶³	Yes	No	N/A	CH ₄ and CO ₂	5% Pt/alumina	2 h with temperature increasing from 200 to 400 °C	2
2004 ⁶⁴	No	Yes	Formic acid (aq)	Nonane-thiol	Ni ^{0b}	6 h at 250 °C and 200 MPa	10–13
2004 ⁵²	No	Yes	Oxalic acid (aq)	Oxalic acid (aq)	Stainless steel	18 h at 300 °C	8–20
2006 ⁴⁸	No	Yes	Formic acid (aq)	Formic acid (aq)	Powdered iron	86 h at 250 °C and 325 bar	2–4
2010 ⁶⁵	Yes	Yes	Water (l) ^c	CO ₂ (aq)	Iron nanoparticles	25–200 h at fixed temp from 80 to 200 °C	1–2
2016 ⁶⁶	Yes	Yes	N/A	Formamide	Meteorite powder	24 h at 140 °C	16, 18
2016 ⁴²	Yes	Yes	N/A	CO (g) and acetylene (g)	NiS	1 week at 105 °C and 2.5 bar	Saturated: 3, 5 Unsaturated: 3, 5, 7, 9
2018 ⁵³	No	Yes	Oxalic acid (g)	Oxalic acid (g)	Stainless steel	>66 h at 175 °C	7–26
2018 ⁶⁷	Yes	Yes	N/A	CO ₂ (g)	Fe ⁰ , Ni ⁰ , Co ⁰	16 h at fixed temp from 30 to 150 °C	1–2
2019 ⁵⁴	Yes	Yes	N/A	HCN, Na ₂ CO ₃ , formaldehyde	Macroporous Ni	Multistep reaction	2–10
2020 ⁴³	No	Yes	Water (l)	CO ₂ (aqueous)	NiS–FeS	50 min at room temperature	1
2020 ⁴⁴	Yes	Yes	H ₂ (g)	CO ₂ (g)	Fe ₃ S ₄ , Fe ₃ O ₄ , Ni ₃ Fe	0–24 h at fixed temp from 60 to 100 °C	1–2

^aUnless otherwise specified, fatty acids were saturated and unbranched. Note: The minimum and maximum length fatty acids may have been synthesized during separate experiments that were reported in the same reference. ^bOther minerals were tested as well, although reactions with Ni⁰ produced linear fatty acids with the greatest number of carbons. ^cH₂ is generated when water oxidizes the Fe.

Table 2. Summary of Electrochemical Experiments That Have Synthesized Fatty Acids^a

year and ref	electric discharge description	carbon source	reaction conditions	number of carbons per fatty acid
1967 ⁷²	1 electrode in solution, 1 electrode in headspace	CH ₄ (g)	96 h of discharge	2–5
1981 ²⁴	1 electrode in solution, 1 electrode in headspace	CH ₄ (g)	24 h of discharge; pH = 8	2–7
2015 ⁷⁸	Both electrodes in solution	CO ₂ (aq)	150 h with electric potential cycling from –0.8 to 0.2 V	1–2
2021 ⁷¹	Both electrodes in headspace	CH ₄ (g)	Discharge alternating on/off for 14 days, pH = 8.7, room temperature	6, 9

^aAll fatty acids were saturated and unbranched.

modern cells.⁶⁰ Ultramafic rocks (relatively Fe- and Mg-rich and Si-poor) are common at some deep sea hydrothermal vents.⁶¹ Could metal-rich minerals within these rocks serve as catalysts for fatty acid synthesis? To address this question, we consider the oxidation state of the mineral surface. Ultramafic mineral surfaces become oxidized by reacting with seawater and generating H₂ in serpentinization reactions.⁶² As noted above, to date, there have been no reports of the synthesis of membrane-forming fatty acids (at least 8 carbons) on oxidized metal surfaces.^{23,45–48} Laboratory experiments failed to produce fatty acids with more than 2 carbons when oxidized olivine (ultramafic mineral with general composition (Fe,Mg)-SiO₄) was the sole catalyst.⁴⁵ In these experiments, the olivine was heated in water for 96 days to allow ample time for oxidation by serpentinization, and formic acid was included as an additional source of H₂.⁴⁵ It is unknown whether olivine surfaces could catalyze fatty acid synthesis before becoming oxidized during serpentinization.

Hydrothermal experiments with oxidation-resistant stainless steel surfaces, clearly not natural settings, do produce fatty acids from formic acid with up to 22 carbons.⁴⁹ However, these experiments also permitted vapor-phase reactions, which may have enabled production of longer fatty acids regardless of the oxidation state of the surface. Thus, in natural hydrothermal settings, it remains unclear if catalytically active, reduced mineral surfaces could persist or be replenished quickly enough to catalyze fatty acid synthesis. Without a suitable catalyst, fatty acid synthesis would likely be slow or nonexistent in hydrothermal environments.⁶³ In conclusion, additional experiments are necessary to determine whether fatty acids could be synthesized in natural hydrothermal settings.

■ ELECTROCHEMICAL SYNTHESIS OF FATTY ACIDS

Electrochemical reactions can generate diverse organic compounds, including amino acids,^{68,69} nucleobases,⁷⁰ and

fatty acids. For example, spark discharges between an electrode in the gaseous headspace and another electrode either in the same headspace⁷¹ or in solution^{24,72} can produce fatty acids. In the experiments summarized in Table 2, CH₄ in the gaseous headspace served as the source of carbon in three of the four experiments generating linear fatty acids of varying lengths with only one experiment reporting chains long enough (at least 8 carbons) to form membranes.⁷¹ An additional electric discharge experiment produced carbon chains of 2–6 carbons with carboxylic acids on both ends.⁷³

The goal of most electrochemical experiments is to simulate lightning strikes through a methane-rich atmosphere on the early Earth. However, attributes of natural lightning strikes are challenging to reproduce in the laboratory. Criado-Reyes et al. used a Tesla coil that generated a 3×10^4 V potential for 7 days at room temperature.⁷¹ In contrast, a natural lightning strike generates a potential of about 10^8 V for <1 s,⁷⁴ and temperatures of the air surrounding a lightning strike can reach 30 000 °C.⁷⁵ Unfortunately, experimental evidence is lacking to describe how fatty acid yields depend on the voltage, duration, or total energy dissipated by laboratory sparking. Chyba and Sagan assumed that electrochemical production of organic molecules on early Earth should depend on the total amount of electrical energy dissipated by lightning strikes and coronal discharges.⁷⁶ Additional experiments are needed to validate this assumption.

Only slightly more is known about the role of the atmosphere and solid surfaces during electrochemical synthesis. Experiments by Schlesinger and Miller have shown that the yield of amino acids during sparking increases with the partial pressure of CH₄;⁷⁷ similar experiments with fatty acids are still needed. Borosilicate glass as a substrate has also been shown to increase the yields of electrochemical fatty acid synthesis.⁷¹ Although borosilicate does not occur naturally, silicates would have been ubiquitous on the early Earth because they are common rock-forming minerals. Additional experiments are necessary to better understand how the yields of electrochemical fatty acid production depend on the gaseous, solid, aqueous, and electrical environments. Nevertheless, the available experimental data suggest that fatty acid synthesis is possible under certain electrochemical conditions.

ALTERNATIVE TYPES OF FATTY ACID SYNTHESIS

In addition to the three main sources reviewed above, there are references in the literature to four alternative types of fatty acid synthesis (Figure 6, Table 3). These include photochemical reactions,^{54,79} irradiation by massive particles,^{80,81} gas-phase ion–molecule reactions,⁸² and redox reactions in aqueous solution.^{83–85}

There have been two reports of fatty acid synthesis by UV photochemistry. By irradiating a gaseous mixture of ethane and ammonia above boiling water (Figure 6A) at 185 and 254 nm, Groth and Weyssenhoff generated fatty acids 1–5 carbons long.⁷⁹ Bonfio et al. used a multistep procedure to synthesize longer-chain fatty acids capable of membrane assembly.⁵⁴ UV irradiation at 254 nm was used in one step, and catalysis by a nickel surface was used in a subsequent step. Photochemistry could provide a plausible explanation for the presence of fatty acids on the surface of the early Earth because UV radiation down to ~200 nm could have penetrated prebiotic atmospheres.⁸⁶ The experiments by Groth and Weyssenhoff did not generate fatty acids when methane was used instead of ethane;⁷⁹ further investigation is needed to determine whether




Figure 6. Alternative reaction types that have demonstrated fatty acid synthesis. (A) Groth and Weyssenhoff used UV photochemistry to convert ethane into fatty acids.⁷⁹ (B) Kaiser et al. irradiated an ultracold (10 K) mixture of ~99% CH₄ and ~1% O₂ with 9 MeV alpha particles to produce fatty acids.⁸⁰ (C) Blagojevic et al. report reactions between gas-phase ions (CH₂⁺ and C₂H₄⁺) and CO to produce fatty acids.⁸² (D) Novotný et al. report the decomposition of monosaccharides into fatty acids under mild alkaline conditions (50 mM NaOH).⁸⁴

radiation from realistic early Earth solar spectra could have enabled photochemical conversion of methane into fatty acids. In contrast, Bonfio et al. used starting materials that are more plausibly prebiotic such as formaldehyde, HCN, NaH₂PO₄, Na₂CO₃, and NaSH,⁵⁴ but these reagents impose a constraint on the geochemical scenario for the early Earth. In a separate experiment by Dworkin et al., membrane-forming amphiphiles were synthesized by irradiating ultracold ices composed of 100:50:1:1 H₂O:CH₃OH:NH₃:CO with 121.6 and ~160 nm UV photons, although the identity of these amphiphiles was not determined.⁸⁷

The second type of “alternative” fatty acid synthesis is irradiation by particles. Experiments of this type were designed to simulate chemistry in the outer solar system, rather than the early Earth. By irradiating an ultracold (10 K) mixture of ~99% CH₄ and ~1% O₂ with 9 MeV alpha particles, Kaiser et al. produced linear fatty acids with 13, 15, or 17 carbons⁸⁰ (Figure 6B). In a subsequent experiment by Kim and Kaiser, an ultracold (10 K) mixture of CO₂ and hydrocarbons (1–6 carbons) was irradiated with 5 keV electrons. The presence of carboxylic acids was confirmed by Fourier-transform infrared spectroscopy, but specific fatty acids were not identified.⁸¹

A third alternative fatty acid synthesis is discussed by Blagojevic et al.⁸² In their experiments, reactions between gas-phase ions (CH₂⁺ and C₂H₄⁺) and molecules (CO) resulted in carboxylic acids with 2 or 3 carbons (Figure 6C). Additional gas-phase ion–molecule reactions to produce formic acid (e.g., CH₄ + O₂⁺ → HCOOH₂⁺ + H) have been suggested but not validated experimentally.⁸⁸ These reactions were suggested to occur in the interstellar medium, and the relevance to early Earth conditions has not been explored.

Table 3. Summary of Experiments That Have Synthesized Fatty Acids by the Mechanisms Illustrated in Figure 6^a

year and ref	reactants	reaction conditions	number of carbons per fatty acid
Photochemistry			
1960 ⁷⁹	Ethane, NH ₃ , H ₂ O	1 week of UV irradiation (185 and 254 nm)	1–3
2019 ⁵⁴	HCN, formaldehyde, Na ₂ CO ₃	Multistep reaction (254 nm UV irradiation and catalysis by Ni surface)	2–10
Irradiation by Massive Particles			
1995 ⁸⁰	CH ₄ , O ₂	Irradiation with 9 MeV alpha particles at 10 K	13, 15, 17
2010 ⁸¹	CO ₂ , hydrocarbons (1–6 carbons)	Irradiation with 5 keV electrons at 10 K	undetermined
Gas-Phase Ion–Molecule Reactions			
2003 ⁸²	CO and either CH ₂ ⁺ or C ₂ H ₄ ⁺	CH ₄ and C ₂ H ₄ are ionized and reacted together in the presence of He and trace H ₂ O at 0.35 Torr and room temperature ^b	2–3
Redox Reactions in Solution			
1963 ⁸³	HCN, NH ₃	90 °C for 18 h	1
2000 ⁸⁵	Fatty aldehydes (either 7 or 8 carbons)	30% H ₂ O ₂ solution at 90 °C for 2 h	7 or 8
2008 ⁸⁴	Glucose, fructose, arabinose, glyceraldehyde, or dihydroxyacetone	50 mM NaOH for 1 h	1–3

^aAll fatty acids were saturated and unbranched. ^bThe authors suggest that their synthesis models low temperature and low pressure environments.

The fourth group of lesser studied fatty acid syntheses are redox reactions. Three sets of redox reactions have been shown to produce fatty acids in solution. The first experiments from Sato et al. use hydrogen peroxide to oxidize fatty aldehydes (7–8 carbons) to produce fatty acids with the same carbon-chain length.⁸⁵ It is unlikely that sufficient hydrogen peroxide would have been present on the early Earth.⁸⁹ In the second set of experiments, Novotný et al. report the decomposition of monosaccharides (3–6 carbons) into linear carboxylic acids (1–3 carbons) under mild alkaline conditions (50 mM NaOH, Figure 6D).⁸⁴ Diverse monosaccharides are obtained in low yield during the formose reaction,⁹⁰ and alkaline lakes on early Earth might have been sites for decomposition into short-chain fatty acids.^{91,92} Finally, Lowe et al. reported production of formic acid and potentially other carboxylic acids by heating a mixture of ammonia and hydrogen cyanide to 90 °C.⁸³ Hydrogen cyanide is considered a prebiotic reagent that can be sequestered and concentrated as ferrocyanide within early Earth environments.^{91,93}

There are numerous additional pathways for the synthesis of short-chain (1–3 carbons) linear fatty acids,⁹⁴ but because membrane assembly requires fatty acids with at least 8 carbons, we have generally omitted them. We are unaware of any abiotic mechanisms by which carbon chains of existing fatty acids are elongated. Modern cells synthesize fatty acids from acetyl-CoA using the sophisticated enzyme complex fatty acid synthetase.⁹⁵ Because the intermediate compounds that are produced during this process are unstable, it is believed that fatty acid synthesis via these reactions would have occurred at a negligible rate on the early Earth before the emergence of enzymes.⁹⁶ Finally, thermodynamic calculations suggest that fatty acids can be synthesized from polyaromatic hydrocarbons,⁹⁷ but to our knowledge, this synthesis has never been demonstrated experimentally.

■ CRITICAL ANALYSIS OF ANALYTICAL TECHNIQUES

Many of the reactions discussed above produce a wide variety of fatty acids and other products. Uniquely identifying products can be challenging, and determining the concentration of each product is even more difficult. In Table 4, we

summarize the reported concentrations of fatty acids from each experiment and comment on potential limitations of the analyses. In general, the papers in Table 4 convincingly identify fatty acids of various lengths but do not provide substantial evidence for the fatty acid concentrations that they report. Moreover, many papers report only relative concentrations of fatty acids (e.g., X% of all products by mass or moles) instead of absolute concentrations (e.g., X moles or X grams), so it is difficult to compare product yields between experiment types. An ideal strategy for the characterization and absolute quantitation of fatty acids was employed by Yuen et al., which involved chromatography and tandem mass spectrometry with isotope-labeled internal standards to eliminate matrix effects.²⁴ Additional synthesis experiments that determine the absolute concentration of fatty acids would be valuable.

■ ESTIMATING THE CONTRIBUTION OF EACH SOURCE TO THE EARLY EARTH FATTY ACID INVENTORY

To gauge the relative importance of each fatty acid source for the origin of cells, we go beyond purely reviewing the literature with a goal of estimating how much fatty acid could be supplied to the early Earth from three sources: delivery by carbonaceous chondrites, catalysis by metal surfaces, and electrochemistry. Many relevant parameters for these estimates lack experimental constraints, so assumptions are needed. Our first assumption is that cells formed during the latter part of the Hadean eon; the full eon was ~4.6 to 4.0 billion years ago (Gya).¹⁸ During the early Hadean, Earth was potentially hit by impactors that were large enough to sterilize the planet's surface and reset the fatty acid inventory, so life must have originated after the last such event.⁹⁸ The median age of estimates for the last ocean-vaporizing impact is ~4.3 Gya,⁹⁹ and we assume that life had originated by 4.0 Gya.¹⁸ In the subsections below, we construct back-of-the-envelope estimates for the total mass of fatty acids supplied to Earth during this interval. Our estimates are based on empirical data from the literature, and we indicate when existing models from the literature are applied. We articulate our assumptions and suggest experiments that will help to refine these estimates. Our estimates cannot distinguish fatty acids that form

Table 4. Summary of Analytical Techniques in Each Report^a

year and ref	carbons per fatty acid	yield information	analytical techniques	identification of products	quantitation of product concentrations
1976 ⁵⁰	10–16	None	GC-MS	Method 1: Metal Surface Catalysis Unambiguous identification by comparing fragmentation spectra from (derivatized) products with fragmentation spectra from authentic standards	N/A
1979 ²³	5–21	≤0.08% yield of normal fatty acids (by mass, relative to initial CO)	GC-MS GC-FID	Unclear how products were identified. Comparisons with authentic standards were not shown.	No data were shown to validate the FID procedure for quantifying product concentrations.
1997 ⁴¹	2	≤40% yield (by moles, relative to initial CH ₃ SH)	GC-MS	Unclear how products were identified. Comparisons with authentic standards were not shown.	No data were shown to validate the GC-MS procedure for quantifying product concentrations.
1999 ⁴⁹	6–22	≤20.8% of all products are fatty acids (by moles)	GC-MS GC-FID	Unambiguous identification of select products by comparing fragmentation spectra and retention times with authentic standards.	No data were shown to validate the procedure for quantifying product concentrations.
2001 ⁵¹	7–18	≤20% yield of fatty acids (by mass, relative to total mass of reaction extract)	GC-MS	Unambiguous identification by comparing fragmentation spectra and retention times with authentic standards.	No data were shown to validate the GC-MS procedure for quantifying product concentrations.
2003 ⁶³	2	None	Diffuse reflectance infrared Fourier transform spectroscopy	Using diffuse reflectance infrared Fourier transform spectroscopy, experimental spectra were compared to authentic standards.	N/A
2004 ⁶⁴	10–13	C10: 53.7% yield	GC-MS	Unambiguous identification by comparing fragmentation spectra with authentic standards.	Calibration curves were described in the text to validate the procedure for quantifying product concentrations. Pentadecane was used as an internal standard for all compounds.
2004 ⁵²	8–20	2.8% of all products are fatty acids (unclear whether by mass or moles)	GC-MS	Unambiguous identification by comparing fragmentation spectra and retention times with authentic standards.	No data were shown to validate the MS procedure for quantifying product concentrations.
2006 ⁴⁸	2–4	C2: 1% yield C3: 0.06% yield C4: <0.02% yield (by moles, relative to initial formic acid)	GC-MS GC-FID	Unclear how products were identified. Comparisons with authentic standards were not shown.	No data were shown to validate the FID procedure for quantifying product concentrations.
2010 ⁶⁵	1–2	C2: 9.0 mM C3: 3.5 mM	GC-MS	Unambiguous identification by comparing fragmentation spectra with authentic standards.	Calibration curves were shown to validate the procedure for quantifying product concentrations. Internal standards were not used.
2016 ⁶⁶	16, 18	C16: 0.011% yield C18: 0.02% yield (by moles, relative to initial NH ₂ CHO)	GC-MS	Unambiguous identification by comparing fragmentation spectra with authentic standards and literature references.	No data were shown to validate the MS procedure for quantifying product concentrations. Betulinic acid was mentioned as the internal standard for all products.
2016 ⁴²	Saturated: 3, 5 Unsaturated: 3, 5, 7, 9 C5 unsaturated: 7.9 mM C7 unsaturated: 0.44 mM C9 unsaturated: 0.011 mM	C3: 7.1 mM C5: 0.3 mM C3 unsaturated: 3.9 mM	GC-MS	For most products, unambiguous identification by comparing fragmentation spectra with authentic standards. For select products, identification was inferred on the basis of mass spectra and relative retention times.	Calibration curves were mentioned (but not shown) to validate the procedure for quantifying product concentrations. For some products, calibration curves were made with authentic standards. For other products, related compounds were used for calibration. Internal standards were not used.

Table 4. continued

year and ref	carbons per fatty acid	yield information	analytical techniques	identification of products	quantitation of product concentrations
2018 ⁵³	7–26	7.8% of all products are fatty acids (unclear whether by mass or moles)	GC-MS	Method 1: Metal Surface Catalysis Unambiguous identification by comparing fragmentation spectra with authentic standards.	No data were shown to validate the GC-MS procedure for quantifying product concentrations. <i>N</i> -Eicosane- <i>d</i> ₄₂ was mentioned as the internal standard for all experiments.
2018 ⁶⁷	1–2	Formic acid: ≤0.21% yield Acetic acid: ≤0.053% yield (by moles, relative to initial CO ₂)	GC-MS ¹ H NMR	Unambiguous identification by comparing GC-MS fragmentation spectra and retention times with authentic standards. In addition, ¹ H NMR spectra were compared with authentic standards.	H NMR calibration curves were shown to validate the procedure for quantifying product concentrations. Authentic standards were used, and DSS-Na was used as an internal standard.
2019 ⁵⁴	2–10	36% of all 8-carbon products are fatty acids (by moles) C1: 1.5 μM	¹ H NMR	Unambiguous identification by comparing ¹ H NMR spectra with authentic standards.	Data from ¹ H NMR spectra were used to determine the relative abundance of fatty acids. Internal standards were not discussed.
2020 ⁴³	1		¹ H NMR 13C-NMR	Unambiguous identification by comparing ¹ H NMR spectra with authentic standards.	H NMR spectra with an internal standard (acetone) were used to quantitate formate concentration. To validate the quantitation, additional formate was spiked in and a corresponding increase in ¹ H NMR signal was observed.
2020 ⁴⁴	1–2	C1: ~20% yield C2: ~0.4% yield (by moles, relative to CO ₂)	LC-MS LC-UV ¹ H NMR	Unambiguous identification by comparing ¹ H NMR spectra with authentic standards.	H NMR calibration curves were mentioned (not shown) to validate the procedure for quantifying product concentrations. Authentic standards were used, and DSS-Na was used as an internal standard.
1967 ⁷²	2–5	C2: 1.2% yield C3: 0.86% yield C4: 0.088% yield C5: 0.066% yield (by moles, relative to initial CH ₄)	GC-MS	Method 2: Electrochemistry Unambiguous identification by comparing fragmentation spectra and retention times with authentic standards.	No data were shown to validate the MS procedure for quantifying product concentrations.
1981 ²⁴	2–7	C2: 0.10% yield C3: 0.68% yield C4: 0.014% yield C5: 0.0050% yield C6: 0.00073% yield C7: 0.00025% yield (by moles, relative to initial CH ₄)	GC-MS	Unambiguous identification by comparing fragmentation spectra and retention times with authentic standards.	Calibration curves were shown to validate the procedure for quantifying product concentrations. Singly deuterated internal standards were used for each product.
2015 ⁷⁸	1–2	C1: ~1 μmoles C2: ~0.5 μmoles	¹ H NMR	Unambiguous identification by comparing ¹ H NMR spectra with authentic standards.	H NMR calibration curves were shown (for formic acid) to validate the procedure for quantifying product concentrations. Authentic standards were used and Me ₂ Si was used as an internal standard.
2021 ⁷¹	6, 9	C6: ≤0.0078% yield C9: ≤0.016% yield (by mass, relative to total mass of reaction extract)	GC-MS	Unambiguous identification by comparing fragmentation spectra and retention times with authentic standards.	No data were shown to validate the GC-MS procedure for quantifying product concentrations.

Table 4. continued

year and ref	carbons per fatty acid	yield information	analytical techniques	identification of products	quantitation of product concentrations
1960 ⁷⁹	1–3	C1: 82 μ moles C2: 234 μ moles Acid–base titration C3: 15 μ moles	Silica gel chromatography	Method 3: Photochemistry Identification of products by comparison of retention times with authentic standards.	Silica gel chromatography was used to separate C1, C2, and C3 products. Each putatively pure product was titrated with NaOH to find the equivalence point.
2019 ⁵⁴	2–10	36% of all 8-carbon products are fatty acids (by moles)	¹ H NMR	Unambiguous identification by comparing ¹ H NMR spectra with authentic standards.	Data from ¹ H NMR spectra were shown and used to determine the relative abundance of fatty acids. Internal standards were not mentioned.
1995 ⁸⁰	13, 15, 17	C13: maximum 18.6 picograms C15: maximum 83.9 picograms C17: maximum 57.0 picograms	GC-MS	Method 4: Irradiation by Massive Particles Unclear how products were identified. Comparison with authentic standards were not mentioned.	No data were shown to validate the MS procedure for quantifying product concentrations.
2010 ⁸¹	Undetermined	10 ¹⁶ molecules/cm ²	FTIR	Regions of the FTIR spectra were considered diagnostic for carboxylic acids: $\nu(\text{O–H})$ stretching = 3500–2500 cm ⁻¹ . $\nu(\text{C=O}) = 1720$ cm ⁻¹ $\nu(\text{C–O}) = 1282$ cm ⁻¹	The intensity of the FTIR spectra at 1720 cm ⁻¹ was used to provide an upper limit on carboxylic acid concentration.
2003 ⁸²	2–3	No data were shown	GC-MS	Method 5: Gas-Phase Ion–Molecule Reactions Unambiguous identification by comparing fragmentation spectra with authentic standards.	N/A
1963 ⁸³	1	No quantitative data were shown	Paper chromatography	Method 6: Redox Reactions in Solution The presence of formic acid was inferred on the basis of the comparison of paper chromatography mobility to an authentic standard, and on the basis of a reaction with ammoniacal silver nitrate.	N/A
2000 ⁸⁵	7 or 8	C7: 73–85% yield C8: 65% yield (unclear whether by mass or moles)	GC-MS	Unambiguous identification by comparing fragmentation spectra with authentic standards.	No data were shown to validate the GC-MS procedure for quantifying product concentrations.
2008 ⁸⁴	1–3	C1: 7–20% yield C2: 0.7–12% yield C3: <0.01–0.2% yield (by moles)	GC-MS	Unambiguous identification by comparing fragmentation spectra with publicly available data and by comparing retention times with authentic standards.	Calibration curves were mentioned (but not shown) to validate the procedure for quantifying product concentrations. An internal standard (heptadecane) was used.

^aUnless otherwise specified, fatty acids are saturated and unbranched.

membranes (more than 8 carbons) from those that do not (less than 8 carbons) because the empirical data that one of our estimates relies on does not do so.

Meteorite Delivery. We estimate the total mass of 2–12 carbon fatty acids delivered to Earth by carbonaceous meteorites, M_s , using the following equation:

$$M_s = M \sum_{L=2}^{12} f_L \quad (2)$$

In eq 2, f_L is the dimensionless fraction of a meteorite's mass that comprises fatty acids of length L , where the length denotes the number of carbons in the fatty acid chain. Values for f_L depend on the type of carbonaceous meteorite, and average values can range from 10^{-9} for 10-carbon fatty acids to 10^{-3} for 2-carbon fatty acids.²⁶ Parameter M is the time-integrated total mass of carbonaceous meteorites with radii of 1–100 m that would impact Earth from 4.3 Gya to 4.0. To estimate M , we integrate the following equation for the mass flux (kg/year) at time t (years) in the past from 4.3 to 4.0 Gya:

$$\dot{m} = 8.9C(1 + 1.6 \times 10^{-10} e^{t/\tau})(m_{\max}^{0.46} - m_{\min}^{0.46})q \quad (3)$$

This equation is adapted from Chyba and Sagan.¹⁰⁰ In eq 3, \dot{m} is the mass flux (mass/year) of the total mass of carbonaceous meteorites with mass from m_{\min} to m_{\max} that would impact Earth per year; C is the frequency of carbonaceous meteorites relative to all types of meteorites (~4%) observed in the meteorite fall record;¹⁰¹ τ is the decay constant of 144 million years for the impactor population; m_{\min} is taken as the mass of a meteorite with a 1 m radius; m_{\max} is taken as the mass of a meteorite with a 100 m radius; q is $1 \text{ kg}^{0.54}/\text{year}$. We assume that all meteorites are spherical with a uniform density (2.1 g/cm^3 for CM-type meteorites³⁸ or 3.2 g/cm^3 for C2-type meteorites¹⁰²), so we can calculate the mass of a meteorite in kilograms from its radius in meters. Meteorites with radii less than 100 m can fragment into pieces that impact the Earth's surface with low enough energy that fatty acids are preserved.^{28,29} Using eq 3, we estimate that $\sim 10^{15}$ kg of carbonaceous meteorites with radii between 1 and 100 m would impact Earth from 4.3 to 4.0 Gya.

Assuming that all carbonaceous meteorites are either CM1-type or C2-type, we use eqs 2 and 3 to calculate bounds for the total mass of fatty acids with a length of 2–12 carbons that could have been delivered to Earth from 4.3 to 4.0 Gya. We find that 10^{10} – 10^{13} kg of fatty acids with a length of 2–12 carbons could have been delivered to Earth from 4.3 to 4.0 Gya by carbonaceous meteorites. Our estimate varies over 3 orders of magnitude because the abundance of fatty acids on each type of meteorite (f_L) varies considerably between different meteorite types. We consider only carbonaceous chondrites because they are the most well-characterized type of meteorite that can deliver fatty acids. We do not quantify additional uncertainties in the meteorite flux, nor do we quantify the influence of meteorite size, impact velocity, and impact angle on the fraction of fatty acids that survive impact.¹⁰³

Catalysis by Metal Surfaces after Vesta-Sized Impact.

Next, we consider fatty acid synthesis on metal surfaces in the wake of a large Vesta-sized ($\sim 10^{20}$ kg) asteroid impact on the early Earth. Such an impact would deliver iron to Earth's surface, which could both act as a catalyst for fatty acid

synthesis and generate a H_2 -rich atmosphere by reactions with steam from the vaporized ocean,⁵⁷ as in eq 4:



An H_2 -rich atmosphere appears to be required for fatty acid synthesis by metal catalysts (Table 1). Such a massive asteroid impact would generate a high enough surface temperature to destroy most organic molecules on Earth; then, as the Earth cooled, the synthesis of fatty acids could occur in the H_2 -rich atmosphere. We consider only the last impactor that would reset Earth's fatty acid inventory, which most likely hit Earth between 4.4 and 4.1 Gya.⁹⁹ Significantly smaller meteorite impacts would not generate high partial pressures of H_2 in the atmosphere because H_2 escapes to space on time scales of $\sim 10^6$ – 10^7 years. Additionally, small impactors would not generate global surface temperatures above $200 \text{ }^\circ\text{C}$, which we assume is the minimum temperature required for fatty acid synthesis. Therefore, smaller impactors would not produce many fatty acids, and we do not consider their contribution here.

To estimate the total mass of fatty acids synthesized by metal catalysts after such an impact, M_c , we assume that the rate of fatty acid synthesis depends linearly on gas pressures according to the following equation:

$$M_c = \int_{t_{400}}^{t_{200}} k_c m_{\text{cat}} P_{\text{H}_2}(t_c) P_{\text{CO}}(t_c) dt_c \quad (5)$$

Here, k_c is an empirical rate constant for fatty acid synthesis, calculated from the results of experiments at $400 \text{ }^\circ\text{C}$ by Nooner and Oro.²³ They provide the only measurement to date of the absolute concentration of fatty acids produced on a metal catalyst in the absence of an aqueous phase. The value of k_c is 7.4×10^{-6} kilograms of 6–18 carbon fatty acids (excluding 12-carbon fatty acids, for which data are not available) per bar of H_2 , per bar of CO , per hour of reaction time, per kilogram of catalytic surface available.²³ Only the summed mass of all fatty acids is reported by Nooner and Oro;²³ there is no information given about the number of moles of individual fatty acids. Here, m_{cat} is the mass of available metal catalyst, in kilograms. P_{H_2} and P_{CO} are the atmospheric partial pressures of hydrogen and carbon monoxide, respectively, in bar; both are functions of time, t_c . The product is integrated over time from t_{400} to t_{200} . t_{400} is the number of hours after the impact until the surface temperature reaches $400 \text{ }^\circ\text{C}$, and t_{200} is the number of hours after the impact until the surface temperature cools to $200 \text{ }^\circ\text{C}$. We assume that fatty acid synthesis occurs at a constant rate k_c when the temperature is within this range and that the reaction does not occur when the temperature is outside this range.

We assume that 33% of the asteroid's mass is iron and that 7% of this iron (i.e., 2.3% of the total asteroid mass) remains available on Earth's surface to catalyze fatty acid synthesis (giving $m_{\text{cat}} = 2.3 \times 10^{18}$ kg) based on a linear extrapolation of the impact simulation performed by Citron and Stewart.⁵⁶ The assumption of $\sim 33\%$ iron mass follows Zahnle et al.,⁵⁵ which is the total iron in high iron enstatite (EH-type) meteorites¹⁰⁴ and also the fraction of Earth's mass in its iron core.¹⁰⁵ Bodies with enstatite composition are candidates for impactors that hit the Earth after the Moon-forming impact and Earth's core formation, although the contribution of carbonaceous versus enstatite compositions is debated.^{106–108} In EH enstatites, which are highly reduced, most of the iron is metallic with a smaller fraction of iron sulfide.¹⁰⁹ In a postimpact vapor plume,

the iron is vaporized into atoms, which condensation sequences show condenses to form metallic iron with subsequent cooling.¹¹⁰

We assume that another fraction of the asteroid's iron (between 1% and 90%) is used to generate H₂ from excess H₂O according to eq 4. We adapted a thermochemical model previously developed by Zahnle et al. to compute P_{H_2} , P_{CO} , and temperature as a function of time after the impact.⁵⁵ Depending on the fraction of the asteroid's iron that reacts with water to produce H₂ gas, we find that P_{H_2} ranges from 10⁻³ to 10⁻¹ bar and P_{CO} ranges from 10⁻³ to 10⁻⁷ bar. We calculate that the Earth's postimpact temperature would be between 200 and 400 °C for ~1000 years, and we assume that fatty acid synthesis occurs at a constant rate during this time. Equation 5 estimates that between 10¹¹ and 10¹⁵ kg of fatty acids with a length 6–18 of carbons (excluding 12 carbons, for which data are not available) would be synthesized using metal catalysts in the wake of a Vesta-size impactor.

Our estimate for the total mass of fatty acids synthesized by metal catalysts is uncertain because many factors in our analysis are poorly constrained. First, it is not known whether reduced metals would remain exposed to the atmosphere on Earth's surface after an extremely massive impact.⁵⁶ Additionally, it is unclear how a H₂O steam atmosphere would influence the rate of fatty acid synthesis; steam can affect the yield of Fischer–Tropsch reactions in different ways, depending on the type of catalyst and the type of reaction product.¹¹¹ Even without steam in the atmosphere, it is uncertain whether the rate of fatty acid synthesis depends linearly on gas partial pressures. Kinetic models for the Fischer–Tropsch process have been developed in industrial settings, which generally do not mimic plausible early Earth conditions; functions for each partial pressure depend on the engineered design of the catalyst.⁵⁸ Additionally, the effect of temperature on the final product distribution also appears to depend on the catalyst design.^{51,112} Separate experiments indicate that the yield of fatty acids depends linearly on the amount of catalyst surface area that is available.⁶⁴ In general, further experiments are necessary to constrain these reaction parameters and the precise functional form for fatty acid production in plausible conditions on the early Earth.

Electrochemical Synthesis. We also construct an estimate for the total mass of fatty acids that could be synthesized by electrochemistry from 4.3 to 4.0 Gya,

$$M_{\text{E}} = RP_{\text{CH}_4}Bt_{\text{E}} \quad (6)$$

where R is an empirical rate constant for the synthesis of 2–7 carbon fatty acids, calculated from electrochemical experiments by Yuen et al.²⁴ The value of R is 1.1×10^{-17} kilograms of 2–7 carbon fatty acids per bar of CH₄, per joule of electrical energy dissipated by sparking, per hour of reaction time.²⁴ Although there are no electrochemical experiments that provide absolute concentrations for membrane-forming (longer than 8 carbons) fatty acids, Yuen et al.²⁴ provide absolute concentrations for the largest number of fatty acids. P_{CH_4} is the partial pressure of methane on the early Earth, which we estimate to be between 10⁻¹⁵ to 10⁻¹ bar after the last Vesta-sized (~10²⁰ kg) asteroid impact. This value depends on the initial preimpact abundance of atmospheric CO₂, the fraction of the impactor's iron that becomes oxidized (1–100%), and the importance of methane-forming catalysts, which may reduce the quench temperature of methane, thereby increasing its abundance.⁵⁵ B is the amount of

electrical energy dissipated by lightning and corona discharges on the Hadean Earth during a year. We assume that B is 1.5×10^{18} joules per year, which is the electrical energy dissipated per year on the modern Earth.⁷⁶ t_{E} is the reaction time in years. We assume t_{E} is 10⁵ years, which is an estimate for the lifetime of a methane-rich atmosphere after a Vesta-sized (~10²⁰ kg) asteroid impact.⁵⁵

By applying these assumptions, we calculate that 10⁻⁴ to 10¹⁰ kg of fatty acids with a length of 2–7 carbons could be synthesized on Earth from 4.3 to 4.0 Gya by electrochemical reactions. Although separate experiments have shown that electrochemical production of membrane forming fatty acids with more than 8 carbons is possible,⁷¹ membrane-forming fatty acids were not detected in the experiments by Yuen et al.,²⁴ so our estimate is not directly informative about membrane formation on early Earth. More data are needed about the absolute concentration of membrane-forming fatty acids produced during electrochemical experiments.

There are uncertainties in our electrochemistry estimate. First, there is insufficient information available to estimate the total energy dissipated by the Tesla coil during the experiments by Yuen et al.²⁴ If we assume that their Tesla coil used 30 000 V potential⁷¹ and 15 A, we can estimate that $\sim 4 \times 10^{10}$ joules of energy were dissipated during their 24 h experiment. Even if we had perfect information about the Tesla coil voltage and current, it might not be straightforward to estimate the amount of energy that is usable for chemical synthesis.¹¹³ Furthermore, we assumed that the yield of fatty acids depends only linearly on the amount of available electrical energy.⁷⁶ Further experiments are necessary to validate this assumption. Experiments by Schlesinger and Miller suggest that the yield of amino acids during sparking does increase linearly with increasing CH₄ partial pressure (below ~ 0.06 bar),⁷⁷ but further experiments are needed to validate this assumption for fatty acids.

Summary of Estimates. We have compared the total mass of fatty acids produced by 3 different sources during the Hadean eon, following the last extremely massive impactor that would have reset Earth's fatty acid inventory. We estimate that 10¹¹ to 10¹⁵ kg of 6–18 carbon fatty acids could have been synthesized by metal catalysts derived from the massive impactor. The total mass of fatty acids that could have been delivered by carbonaceous meteorites is 10¹⁰ to 10¹³ kg of 2–12 carbon fatty acids. The yield of 2–7 carbon fatty acids from electrochemical processes is potentially smaller, between 10⁻⁴ and 10¹⁰ kg. Consequently, an integrated supply of fatty acids to the Earth's surface from all sources (dominated by metal surface production) between $\sim 10^{-4}$ and $\sim 10^0$ kg/m² is possible, given the Earth's surface area of 5.1×10^{14} m².

Ultimately, the local concentration of fatty acids determines whether or not membranes form, so the possible sources should be evaluated by this criterion. Although meteorites could have delivered a significant mass of fatty acids across the Earth's surface, the aqueous concentration of fatty acids in a single waterbody would not have been high enough to form membranes without evaporating a significant volume of water (Figure 3). In contrast, a local stockpile of fatty acids could have been produced on atmosphere-exposed metal surfaces after an extremely massive impact, and subsequent dissolution into water could have allowed membrane formation. Although little is known about the electrochemical synthesis rate for membrane-forming fatty acids, repeated lightning strikes into the same small waterbody seem unlikely, so it is unclear

whether a high enough local concentration of membrane-forming fatty acids could have formed via electrochemistry. Fatty acids in aqueous solution can be degraded via photochemistry,¹¹⁴ so fatty acids that are slowly synthesized by electrochemistry may not have attained high enough concentrations to form membranes, whereas a large stockpile of fatty acids dissolving off metal surfaces may have been less sensitive to photochemical degradation.

Although the estimates above have many uncertainties, they are valuable as a first attempt to quantitatively compare fatty acid sources on the early Earth. We hope that future experiments can further constrain these estimates.

ALTERNATIVE AMPHIPHILES

In addition to fatty acids, alternative types of amphiphiles may have been synthesized on the early Earth,^{115–117} and these amphiphiles might have incorporated into the membranes of the earliest cells. For example, alcohols are commonly produced along with fatty acids in many experiments,^{48,49,51–54} and long-chain fatty alcohols are known to stabilize fatty acid membranes.^{5,118} An excess of long-chain fatty alcohols form oil droplets, and the oil may disrupt membranes. In addition, phase-separated coacervates could have served as another type of prebiotic compartment,^{119,120} and fatty acid membranes may have even assembled around such coacervate compartments.¹²¹

CONCLUSIONS

Fatty acids can assemble into membranes and could have formed the boundaries for the first cells. Our Review highlights multiple potential sources of fatty acids on the early Earth. The three most well-characterized sources are meteorite delivery, synthesis on metal surfaces, and synthesis by electrochemistry. Other reactions involving photochemistry, irradiation by massive particles, ion–molecule reactions, and diverse redox reactions in aqueous solution may have also produced fatty acids in natural environments. To refine quantitative estimates for the relative importance of each fatty acid source, more detailed constraints are needed. We highlight the following questions to help any future experiments have the widest possible impact:

- How do the yields of fatty acids from metal-catalyzed reactions depend on the temperature and the partial pressures of gases such as H₂, CO₂, and H₂O? Data would help to constrain estimates for fatty acid production following ocean-vaporizing impacts.
- Can membrane-forming fatty acids (>8 carbons) form in hydrothermal experiments with nonoxidized, ultramafic minerals as the sole catalyst? Data would elucidate the potential for fatty acid production at hydrothermal vents.
- How do the yields of fatty acids depend on the voltage, duration, or total energy dissipated during electrical sparking? Data would help to constrain estimates for fatty acid production during lightning strikes.
- What are the absolute concentrations of fatty acids in these types of experiments?

In summary, our analysis suggests that fatty acids could have been available on the early Earth. We have not assessed whether those fatty acids would have been sufficiently concentrated to assemble into membranes except in the limited case of small meteorite fragments delivered into aqueous environments. For fatty acids supplied via alternative

sources, further data are required to assess the potential for membrane formation. By investigating possible sources of fatty acids on the early Earth, we hope to constrain the environmental setting for the origin of cells.

AUTHOR INFORMATION

Corresponding Authors

Zachary R. Cohen – Department of Chemistry and Astrobiology Program, University of Washington, Seattle, Washington 98195, United States; orcid.org/0000-0002-1257-623X; Email: zrcohen2@uw.edu

David C. Catling – Department of Earth and Space Sciences and Astrobiology Program, University of Washington, Seattle, Washington 98195, United States; orcid.org/0000-0001-5646-120X; Email: dcating@uw.edu

Authors

Zoe R. Todd – Department of Earth and Space Sciences, University of Washington, Seattle, Washington 98195, United States; orcid.org/0000-0001-6116-4285

Nicholas Wogan – Department of Earth and Space Sciences and Astrobiology Program, University of Washington, Seattle, Washington 98195, United States

Roy A. Black – Department of Chemistry, University of Washington, Seattle, Washington 98195, United States; orcid.org/0000-0002-3914-5747

Sarah L. Keller – Department of Chemistry and Astrobiology Program, University of Washington, Seattle, Washington 98195, United States; orcid.org/0000-0001-5919-0185

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acsearthspacechem.2c00168>

Author Contributions

Z.R.C. conducted the literature review. Z.R.C., Z.R.T., and N.W. calculated fatty acid abundances. Z.R.C., Z.R.T., N.W., R.A.B, S.L.K, and D.C.C. wrote the paper.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank Milomir Suvira for helpful discussions of electrochemical sparking, Martin Sadilek for helpful discussions of analytical methods, Kevin Zahnle for helpful comments on all portions of the manuscript, and Ben K. D. Pearce for sharing raw data on meteoritic fatty acid abundances that he and his colleagues compiled.²⁶ This work was supported in part by grant NNX17AK86G (Exobiology) from NASA to S.L.K., by a grant (MCB 1925731) from the NSF to S.L.K., and by a grant (511570FY20, DCC) from the Simons Foundation to D.C.C. Z.R.C. was funded by an NSF fellowship (NSF GRFP DGE 1762114).

ABBREVIATIONS

General

Gya, billion years ago

C_n, an organic molecule containing “n” carbons

°C, degrees Celsius

K, degrees Kelvin

h, hours

UV, ultraviolet

mM, millimolar

ppb, parts per billion

ppm, parts per million
 m, meter
 km, kilometer
 cm, centimeter
 nm, nanometer
 kg, kilogram
 V, volt
 s, second
 MeV, mega electronvolt
 KeV, kilo electronvolt
 pH, negative log₁₀ of proton concentration
 pK_a, negative log₁₀ of the equilibrium constant (K_a) for dissociation into a proton and the conjugate base

Analytical techniques

GC, gas chromatography
 LC, liquid chromatography
 MS, mass spectrometry
 FID, flame ionization detector
¹H-NMR, proton nuclear magnetic resonance spectroscopy
¹³C-NMR, carbon-13 nuclear magnetic resonance spectroscopy
 FTIR, Fourier-transform infrared spectroscopy

Chemical names

H₂O, water
 H₂, hydrogen gas
 D₂, deuterium gas
 CO₂, carbon dioxide
 CO, carbon monoxide
 CH₄, methane
 O₂, oxygen gas
 NH₃, ammonia
 He, helium
 HCN, hydrogen cyanide
 Na₂CO₃, sodium carbonate
 NaH₂PO₄, monosodium phosphate
 NaSH, sodium hydrosulfide
 K₂CO₃, potassium carbonate
 CH₃OH, methanol
 NaOH, sodium hydroxide
 H₂O₂, hydrogen peroxide
 Ni, nickel
 Fe, iron
 Mg, magnesium
 Si, silicon
 Pt, platinum
 Co, cobalt
 FeS, iron sulfide
 NiS, nickel sulfide
 FeO, iron (II) oxide
 Fe₃O₄, iron (II, III) oxide
 CH₃SH, methanethiol
 CoA, coenzyme A

REFERENCES

- Deamer, D.; Dworkin, J. P.; Sandford, S. A.; Bernstein, M. P.; Allamandola, L. J. The First Cell Membranes. *Astrobiology* **2002**, *2* (4), 371–381.
- Black, R. A.; Blosser, M. C.; Stottrup, B. L.; Tavakley, R.; Deamer, D. W.; Keller, S. L. Nucleobases Bind to and Stabilize Aggregates of a Prebiotic Amphiphile, Providing a Viable Mechanism for the Emergence of Protocells. *Proc. Natl. Acad. Sci. USA* **2013**, *110* (33), 13272.
- Cornell, C. E.; Black, R. A.; Xue, M.; Litz, H. E.; Ramsay, A.; Gordon, M.; Mileant, A.; Cohen, Z. R.; Williams, J. A.; Lee, K. K.; Drobny, G. P.; Keller, S. L. Prebiotic Amino Acids Bind to and Stabilize Prebiotic Fatty Acid Membranes. *Proc. Natl. Acad. Sci. USA* **2019**, *116* (35), 17239–17244.
- Xue, M.; Black, R. A.; Cohen, Z. R.; Roehrich, A.; Drobny, G. P.; Keller, S. L. Binding of Dipeptides to Fatty Acid Membranes Explains Their Colocalization in Protocells but Does Not Select for Them Relative to Unjoined Amino Acids. *J. Phys. Chem. B* **2021**, *125* (29), 7933–7939.
- Apel, C. L.; Deamer, D. W.; Mautner, M. N. Self-Assembled Vesicles of Monocarboxylic Acids and Alcohols: Conditions for Stability and for the Encapsulation of Biopolymers. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **2002**, *1559* (1), 1–9.
- Cistola, D. P.; Hamilton, J. A.; Jackson, D.; Small, D. M. Ionization and Phase Behavior of Fatty Acids in Water: Application of the Gibbs Phase Rule. *Biochemistry* **1988**, *27* (6), 1881–1888.
- Jordan, S. F.; Ramm, H.; Zheludev, I. N.; Hartley, A. M.; Maréchal, A.; Lane, N. Promotion of Protocell Self-Assembly from Mixed Amphiphiles at the Origin of Life. *Nat. Ecol. Evol.* **2019**, *3*, 1705.
- Caschera, F.; de la Serna, J. B.; Löffler, P. M. G.; Rasmussen, T. E.; Hanczyc, M. M.; Bagatolli, L. A.; Monnard, P.-A. Stable Vesicles Composed of Monocarboxylic or Dicarboxylic Fatty Acids and Trimethylammonium Amphiphiles. *Langmuir* **2011**, *27* (23), 14078–14090.
- Chen, I. A.; Walde, P. From Self-Assembled Vesicles to Protocells. *Cold Spring Harb Perspect Biol.* **2010**, *2* (7), a002170.
- Mansy, S. S.; Schrum, J. P.; Krishnamurthy, M.; Tobé, S.; Treco, D. A.; Szostak, J. W. Template-Directed Synthesis of a Genetic Polymer in a Model Protocell. *Nature* **2008**, *454* (7200), 122–125.
- Chen, I. A.; Szostak, J. W. A Kinetic Study of the Growth of Fatty Acid Vesicles. *Biophys. J.* **2004**, *87* (2), 988–998.
- Chen, I. A.; Roberts, R. W.; Szostak, J. W. The Emergence of Competition Between Model Protocells. *Science* **2004**, *305* (5689), 1474–1476.
- Zhu, T. F.; Szostak, J. W. Coupled Growth and Division of Model Protocell Membranes. *J. Am. Chem. Soc.* **2009**, *131* (15), 5705–5713.
- Toparal, Ö. D.; Wang, A.; Mansy, S. S. Population-Level Membrane Diversity Triggers Growth and Division of Protocells. *JACS Au* **2021**, *1* (5), 560–568.
- Hanczyc, M. M.; Fujikawa, S. M.; Szostak, J. W. Experimental Models of Primitive Cellular Compartments: Encapsulation, Growth, and Division. *Science* **2003**, *302* (5645), 618–622.
- Coreta-Gomes, F. M.; Vaz, W. L. C.; Moreno, M. J. Effect of Acyl Chain Length on the Rate of Phospholipid Flip-Flop and Intermembrane Transfer. *J. Membr. Biol.* **2018**, *251* (3), 431–442.
- Budin, I.; Szostak, J. W. Physical Effects Underlying the Transition from Primitive to Modern Cell Membranes. *Proc. Natl. Acad. Sci. USA* **2011**, *108* (13), 5249–5254.
- Bell, E. A.; Boehnke, P.; Harrison, T. M.; Mao, W. L. Potentially Biogenic Carbon Preserved in a 4.1 Billion-Year-Old Zircon. *Proc. Natl. Acad. Sci. USA* **2015**, *112* (47), 14518.
- Lawless, J. G.; Yuen, G. U. Quantification of Monocarboxylic Acids in the Murchison Carbonaceous Meteorite. *Nature* **1979**, *282* (5737), 396–398.
- Naraoka, H.; Shimoyama, A.; Harada, K. Molecular Distribution of Monocarboxylic Acids in Asuka Carbonaceous Chondrites from Antarctica. *Orig. Life Evol. Biosph.* **1999**, *29* (2), 187–201.
- Aponte, J. C.; Alexandre, M. R.; Wang, Y.; Brearley, A. J.; Alexander, C. M. O.; Huang, Y. Effects of Secondary Alteration on the Composition of Free and IOM-Derived Monocarboxylic Acids in Carbonaceous Chondrites. *Geochim. Cosmochim. Acta* **2011**, *75* (9), 2309–2323.
- Hilts, R. W.; Herd, C. D. K.; Simkus, D. N.; Slater, G. F. Soluble Organic Compounds in the Tagish Lake Meteorite. *Meteoritics & Planetary Science* **2014**, *49* (4), 526–549.

- (23) Nooner, D. W.; Oro, J. Synthesis of Fatty Acids by a Closed System Fischer–Tropsch Process. In *Hydrocarbon Synthesis from Carbon Monoxide and Hydrogen*; Advances in Chemistry; American Chemical Society, 1979; Vol. 178, pp 159–171.
- (24) Yuen, G. U.; Lawless, J. G.; Edelson, E. H. Quantification of Monocarboxylic Acids from a Spark Discharge Synthesis. *J. Mol. Evol.* **1981**, *17* (1), 43–47.
- (25) Deamer, D. W. Boundary Structures Are Formed by Organic Components of the Murchison Carbonaceous Chondrite. *Nature* **1985**, *317* (6040), 792–794.
- (26) Lai, J. C.-Y.; Pearce, B. K. D.; Pudritz, R. E.; Lee, D. Meteoritic Abundances of Fatty Acids and Potential Reaction Pathways in Planetesimals. *Icarus* **2019**, *319*, 685–700.
- (27) Sephton, M. A. Organic Compounds in Carbonaceous Meteorites. *Nat. Prod. Rep.* **2002**, *19* (3), 292–311.
- (28) Chyba, C. F.; Thomas, P. J.; Brookshaw, L.; Sagan, C. Cometary Delivery of Organic Molecules to the Early Earth. *Science* **1990**, *249* (4967), 366–373.
- (29) Pearce, B. K. D.; Pudritz, R. E.; Semenov, D. A.; Henning, T. K. Origin of the RNA World: The Fate of Nucleobases in Warm Little Ponds. *Proc. Natl. Acad. Sci. USA* **2017**, *114* (43), 11327–11332.
- (30) Zamoždra, S. N.; Kocherov, A. V. Underwater Excavations and Welcoming the Big Meteorite. In *Chelyabinsk Superbolide*; Gorkavii, N., Dudorov, A., Taskaev, S., Eds.; Springer Praxis Books; Springer International Publishing: Cham, 2019; pp 122–147.
- (31) Popova, O. P.; Jenniskens, P.; Emel'yanenko, V.; Kartashova, A.; Biryukov, E.; Khaibrakhmanov, S.; Shuvalov, V.; Rybnov, Y.; Dudorov, A.; Grokhovskiy, V. I.; Badyukov, D. D.; Yin, Q.-Z.; Gural, P. S.; Albers, J.; Granvik, M.; Evers, L. G.; Kuiper, J.; Kharlamov, V.; Solovyov, A.; Rusakov, Y. S.; Korotkiy, S.; Serdyuk, I.; Korochantsev, A. V.; Larionov, M. Yu.; Glazachev, D.; Mayer, A. E.; Gisler, G.; Gladkovsky, S. V.; Wimpenny, J.; Sanborn, M. E.; Yamakawa, A.; Verosub, K. L.; Rowland, D. J.; Roeske, S.; Botto, N. W.; Friedrich, J. M.; Zolensky, M. E.; Le, L.; Ross, D.; Ziegler, K.; Nakamura, T.; Ahn, I.; Lee, J. I.; Zhou, Q.; Li, X.-H.; Li, Q.-L.; Liu, Y.; Tang, G.-Q.; Hiroi, T.; Sears, D.; Weinstein, I. A.; Vokhmintsev, A. S.; Ishchenko, A. V.; Schmitt-Kopplin, P.; Hertkorn, N.; Nagao, K.; Haba, M. K.; Komatsu, M.; Mikouchi, T.; the Chelyabinsk Airburst Consortium. Chelyabinsk Airburst, Damage Assessment, Meteorite Recovery, and Characterization. *Science* **2013**, *342* (6162), 1069–1073.
- (32) Artemieva, N. A.; Shuvalov, V. V. From Tunguska to Chelyabinsk via Jupiter. *Annual Review of Earth and Planetary Sciences* **2016**, *44* (1), 37–56.
- (33) Mehta, C.; Perez, A.; Thompson, G.; Pasek, M. A. Caveats to Exogenous Organic Delivery from Ablation, Dilution, and Thermal Degradation. *Life* **2018**, *8* (2), 13.
- (34) Clark, B. C.; Kolb, V. M. Comet Pond II: Synergistic Intersection of Concentrated Extraterrestrial Materials and Planetary Environments to Form Procreative Darwinian Ponds. *Life* **2018**, *8* (2), 12.
- (35) Cape, J. L.; Monnard, P.-A.; Boncella, J. M. Prebiotically Relevant Mixed Fatty Acid Vesicles Support Anionic Solute Encapsulation and Photochemically Catalyzed Trans-Membrane Charge Transport. *Chem. Sci.* **2011**, *2* (4), 661–671.
- (36) Budin, I.; Prywes, N.; Zhang, N.; Szostak, J. W. Chain-Length Heterogeneity Allows for the Assembly of Fatty Acid Vesicles in Dilute Solutions. *Biophys. J.* **2014**, *107* (7), 1582–1590.
- (37) Trigo-Rodríguez, J. M.; Rimola, A.; Tanbakouei, S.; Soto, V. C.; Lee, M. Accretion of Water in Carbonaceous Chondrites: Current Evidence and Implications for the Delivery of Water to Early Earth. *Space Sci. Rev.* **2019**, *215* (1), 18.
- (38) Britt, D. T.; Consolmagno, G. J. S. Stony Meteorite Porosities and Densities: A Review of the Data through 2001. *Meteoritics & Planetary Science* **2003**, *38* (8), 1161–1180.
- (39) de Klerk, A. Fischer–Tropsch Synthesis. In *Fischer–Tropsch Refining*; John Wiley & Sons, Ltd.: Hoboken, NJ; pp 73–103.
- (40) Schulz, H.; Beck, K.; Erich, E. Mechanism of the Fischer–Tropsch Process. In *Studies in Surface Science and Catalysis*; Bibby, D. M., Chang, C. D., Howe, R. F., Yurchak, S., Eds.; Methane Conversion; Elsevier, 1988; Vol. 36, pp 457–471.
- (41) Huber, C.; Wächtershäuser, G. Activated Acetic Acid by Carbon Fixation on (Fe,Ni)S Under Primordial Conditions. *Science* **1997**, *276* (5310), 245–247.
- (42) Scheidler, C.; Sobotta, J.; Eisenreich, W.; Wächtershäuser, G.; Huber, C. Unsaturated C₃,₅,₇,₉-Monocarboxylic Acids by Aqueous, One-Pot Carbon Fixation: Possible Relevance for the Origin of Life. *Sci. Rep.* **2016**, *6* (1), 1–7.
- (43) Hudson, R.; de Graaf, R.; Strandoo Rodin, M.; Ohno, A.; Lane, N.; McGlynn, S. E.; Yamada, Y. M. A.; Nakamura, R.; Barge, L. M.; Braun, D.; Sojo, V. CO₂ Reduction Driven by a PH Gradient. *Proc. Natl. Acad. Sci. USA* **2020**, *117* (37), 22873–22879.
- (44) Preiner, M.; Igarashi, K.; Muchowska, K. B.; Yu, M.; Varma, S. J.; Kleinermaans, K.; Nobu, M. K.; Kamagata, Y.; Tüysüz, H.; Moran, J.; Martin, W. F. A Hydrogen-Dependent Geochemical Analogue of Primordial Carbon and Energy Metabolism. *Nat. Ecol. Evol.* **2020**, *4* (4), 534–542.
- (45) McCollom, T. M.; Seewald, J. S. A Reassessment of the Potential for Reduction of Dissolved CO₂ to Hydrocarbons during Serpentinization of Olivine. *Geochim. Cosmochim. Acta* **2001**, *65* (21), 3769–3778.
- (46) McCollom, T. M.; Seewald, J. S. Experimental Constraints on the Hydrothermal Reactivity of Organic Acids and Acid Anions: I. Formic Acid and Formate. *Geochim. Cosmochim. Acta* **2003**, *67* (19), 3625–3644.
- (47) McCollom, T. M.; Seewald, J. S. Experimental Study of the Hydrothermal Reactivity of Organic Acids and Acid Anions: II. Acetic Acid, Acetate, and Valeric Acid. *Geochim. Cosmochim. Acta* **2003**, *67* (19), 3645–3664.
- (48) McCollom, T. M.; Seewald, J. S. Carbon Isotope Composition of Organic Compounds Produced by Abiotic Synthesis under Hydrothermal Conditions. *Earth and Planetary Science Letters* **2006**, *243* (1), 74–84.
- (49) McCollom, T. M.; Ritter, G.; Simoneit, B. R. T. Lipid Synthesis Under Hydrothermal Conditions by Fischer–Tropsch-Type Reactions. *Orig. Life Evol. Biosph.* **1999**, *29* (2), 153–166.
- (50) Nooner, D. W.; Gibert, J. M.; Gelpi, E.; Oró, J. Closed System Fischer–Tropsch Synthesis over Meteoritic Iron, Iron Ore and Nickel-Iron Alloy. *Geochim. Cosmochim. Acta* **1976**, *40* (8), 915–924.
- (51) Rushdi, A. I.; Simoneit, B. R. T. Lipid Formation by Aqueous Fischer–Tropsch-Type Synthesis over a Temperature Range of 100 to 400 °C. *Orig. Life Evol. Biosph.* **2001**, *31* (1), 103–118.
- (52) Rushdi, A. I.; Simoneit, B. R. T. Condensation Reactions and Formation of Amides, Esters, and Nitriles Under Hydrothermal Conditions. *Astrobiology* **2004**, *4* (2), 211–224.
- (53) Mißbach, H.; Schmidt, B. C.; Duda, J.-P.; Lünsdorf, N. K.; Goetz, W.; Thiel, V. Assessing the Diversity of Lipids Formed via Fischer–Tropsch-Type Reactions. *Org. Geochem.* **2018**, *119*, 110–121.
- (54) Bonfio, C.; Caumes, C.; Duffy, C. D.; Patel, B. H.; Percivalle, C.; Tsanakopoulou, M.; Sutherland, J. D. Length-Selective Synthesis of Acylglycerol-Phosphates through Energy-Dissipative Cycling. *J. Am. Chem. Soc.* **2019**, *141* (9), 3934–3939.
- (55) Zahnle, K. J.; Lupu, R.; Catling, D. C.; Wogan, N. Creation and Evolution of Impact-Generated Reduced Atmospheres of Early Earth. *Planet. Sci. J.* **2020**, *1* (1), 11.
- (56) Citron, R. I.; Stewart, S. T. Large Impacts onto the Early Earth: Planetary Sterilization and Iron Delivery. *arXiv* **2022**, arXiv:2201.09349; <https://arxiv.org/abs/2201.09349>.
- (57) Benner, S. A.; Bell, E. A.; Biondi, E.; Brassler, R.; Carell, T.; Kim, H.-J.; Mojzsis, S. J.; Omran, A.; Pasek, M. A.; Trail, D. When Did Life Likely Emerge on Earth in an RNA-First Process? *ChemSystemsChem.* **2020**, *2* (2), e1900035.
- (58) Zhou, L.-P.; Hao, X.; Gao, J.-H.; Yang, Y.; Wu, B.-S.; Xu, J.; Xu, Y.-Y.; Li, Y.-W. Studies and Discriminations of the Kinetic Models for the Iron-Based Fischer–Tropsch Catalytic Reaction in a Recycle Slurry Reactor. *Energy Fuels* **2011**, *25* (1), 52–59.

- (59) Shock, E.; Canovas, P. The Potential for Abiotic Organic Synthesis and Biosynthesis at Seafloor Hydrothermal Systems. *Geofluids* **2011**, *10* (1–2), 161–192.
- (60) Kohn, C.; Charlou, J. L.; Donval, J. P.; Holm, N. G.; Dehairs, F.; Bouillon, S. Hydrocarbons and Oxidized Organic Compounds in Hydrothermal Fluids from Rainbow and Lost City Ultramafic-Hosted Vents. *Chem. Geol.* **2009**, *258* (3), 299–314.
- (61) Bach, W.; Banerjee, N. R.; Dick, H. J. B.; Baker, E. T. Discovery of Ancient and Active Hydrothermal Systems along the Ultra-Slow Spreading Southwest Indian Ridge 10°–16°E. *Geochemistry, Geophysics, Geosystems* **2002**, *3* (7), 1–14.
- (62) McCollom, T. M.; Seewald, J. S. Abiotic Synthesis of Organic Compounds in Deep-Sea Hydrothermal Environments. *Chem. Rev.* **2007**, *107* (2), 382–401.
- (63) Wilcox, E. M.; Roberts, G. W.; Spivey, J. J. Direct Catalytic Formation of Acetic Acid from CO₂ and Methane. *Catal. Today* **2003**, *88* (1), 83–90.
- (64) Cody, G. D.; Boctor, N. Z.; Brandes, J. A.; Filley, T. R.; Hazen, R. M.; Yoder, H. S. Assaying the Catalytic Potential of Transition Metal Sulfides for Abiotic Carbon Fixation 1 Associate Editor: P. A. O'Day. *Geochim. Cosmochim. Acta* **2004**, *68* (10), 2185–2196.
- (65) He, C.; Tian, G.; Liu, Z.; Feng, S. A Mild Hydrothermal Route to Fix Carbon Dioxide to Simple Carboxylic Acids. *Org. Lett.* **2010**, *12* (4), 649–651.
- (66) Rotelli, L.; Trigo-Rodríguez, J. M.; Moyano-Camero, C. E.; Carota, E.; Botta, L.; Di Mauro, E.; Saladino, R. The Key Role of Meteorites in the Formation of Relevant Prebiotic Molecules in a Formamide/Water Environment. *Sci. Rep.* **2016**, *6* (1), 38888.
- (67) Varma, S. J.; Muchowska, K. B.; Chatelain, P.; Moran, J. Native Iron Reduces CO₂ to Intermediates and End-Products of the Acetyl-CoA Pathway. *Nat. Ecol. Evol.* **2018**, *2* (6), 1019–1024.
- (68) Miller, S. L. A Production of Amino Acids Under Possible Primitive Earth Conditions. *Science* **1953**, *117* (3046), 528–529.
- (69) Johnson, A. P.; Cleaves, H. J.; Dworkin, J. P.; Glavin, D. P.; Lazcano, A.; Bada, J. L. The Miller Volcanic Spark Discharge Experiment. *Science* **2008**, *322* (5900), 404–404.
- (70) Ferus, M.; Pietrucci, F.; Saitta, A. M.; Knížek, A.; Kubelík, P.; Ivanek, O.; Shestivska, V.; Civiš, S. Formation of Nucleobases in a Miller–Urey Reducing Atmosphere. *Proc. Natl. Acad. Sci. USA* **2017**, *114* (17), 4306–4311.
- (71) Criado-Reyes, J.; Bizzarri, B. M.; García-Ruiz, J. M.; Saladino, R.; Di Mauro, E. The Role of Borosilicate Glass in Miller–Urey Experiment. *Sci. Rep.* **2021**, *11* (1), 21009.
- (72) Allen, W. V.; Ponnampertuma, C. A Possible Prebiotic Synthesis of Monocarboxylic Acids. *Biosystems* **1967**, *1* (1), 24–28.
- (73) Zeitman, B.; Chang, S.; Lawless, J. G. Dicarboxylic Acids from Electric Discharge. *Nature* **1974**, *251* (5470), 42–43.
- (74) Williams, E. R. The Electrification of Thunderstorms. *SCIENTIFIC AMERICAN* **1988**, *7*, 13.
- (75) Orville, R. E. A High-Speed Time-Resolved Spectroscopic Study of the Lightning Return Stroke: Part II. A Quantitative Analysis. *Journal of the Atmospheric Sciences* **1968**, *25* (5), 839–851.
- (76) Chyba, C.; Sagan, C. Electrical Energy Sources for Organic Synthesis on the Early Earth. *Origins Life Evol. Biosphere* **1991**, *21* (1), 3–17.
- (77) Schlesinger, G.; Miller, S. L. Prebiotic Synthesis in Atmospheres Containing CH₄, CO, and CO₂. *J. Mol. Evol.* **1983**, *19* (5), 376–382.
- (78) Roldan, A.; Hollingsworth, N.; Roffey, A.; Islam, H.-U.; Goodall, J. B. M.; Catlow, C. R. A.; Darr, J. A.; Bras, W.; Sankar, G.; Holt, K. B.; Hogarth, G.; de Leeuw, N. H. Bio-Inspired CO₂ Conversion by Iron Sulfide Catalysts under Sustainable Conditions. *Chem. Commun.* **2015**, *51* (35), 7501–7504.
- (79) Groth, W. E.; Weyssenhoff, H. v. Photochemical Formation of Organic Compounds from Mixtures of Simple Gases. *Planetary and Space Science* **1960**, *2* (2), 79–85.
- (80) Kaiser, R. I.; Gabrysch, A.; Roessler, K. Cosmic Ray Simulator: A Versatile Apparatus for Quantitative Studies on the Interaction of Cosmic Rays with Frozen Solids by on Line and in Situ Quadrupole Mass Spectrometry and Fourier Transform Infrared Spectroscopy. *Rev. Sci. Instrum.* **1995**, *66* (4), 3058–3066.
- (81) Kim, Y. S.; Kaiser, R. I. Abiotic Formation of Carboxylic Acids (RCOOH) in Interstellar and Solar System Model Ices. *ApJ.* **2010**, *725* (1), 1002–1010.
- (82) Blagojevic, V.; Petrie, S.; Bohme, D. K. Gas-Phase Syntheses for Interstellar Carboxylic and Amino Acids. *Mon. Not. R. Astron. Soc.* **2003**, *339* (1), L7–L11.
- (83) Lowe, C. U.; Rees, M. W.; Markham, R. Synthesis of Complex Organic Compounds from Simple Precursors: Formation of Amino-Acids, Amino-Acid Polymers, Fatty Acids and Purines from Ammonium Cyanide. *Nature* **1963**, *199* (4890), 219–222.
- (84) Novotný, O.; Cejpek, K.; Velíšek, J. Formation of Carboxylic Acids during Degradation of Monosaccharides. *Czech J. Food Sci.* **2008**, *26* (2), 113–131.
- (85) Sato, K.; Hyodo, M.; Takagi, J.; Aoki, M.; Noyori, R. Hydrogen Peroxide Oxidation of Aldehydes to Carboxylic Acids: An Organic Solvent-, Halide- and Metal-Free Procedure. *Tetrahedron Lett.* **2000**, *41* (9), 1439–1442.
- (86) Ranjan, S.; Sasselov, D. D. Constraints on the Early Terrestrial Surface UV Environment Relevant to Prebiotic Chemistry. *Astrobiology* **2017**, *17* (3), 169–204.
- (87) Dworkin, J. P.; Deamer, D. W.; Sandford, S. A.; Allamandola, L. J. Self-Assembling Amphiphilic Molecules: Synthesis in Simulated Interstellar/Precometary Ices. *Proc. Natl. Acad. Sci. USA* **2001**, *98* (3), 815–819.
- (88) Ehrenfreund, P.; d'Hendecourt, L.; Charnley, S.; Ruitkamp, R. Energetic and Thermal Processing of Interstellar Ices. *Journal of Geophysical Research: Planets* **2001**, *106* (E12), 33291–33301.
- (89) Pecoits, E.; Smith, M. L.; Catling, D. C.; Philippot, P.; Kappler, A.; Konhauser, K. O. Atmospheric Hydrogen Peroxide and Eoarchean Iron Formations. *Geobiology* **2015**, *13* (1), 1–14.
- (90) Omran, A.; Menor-Salvan, C.; Springsteen, G.; Pasek, M. The Messy Alkaline Formose Reaction and Its Link to Metabolism. *Life* **2020**, *10* (8), 125.
- (91) Toner, J. D.; Catling, D. C. Alkaline Lake Settings for Concentrated Prebiotic Cyanide and the Origin of Life. *Geochim. Cosmochim. Acta* **2019**, *260*, 124–132.
- (92) Toner, J. D.; Catling, D. C. A Carbonate-Rich Lake Solution to the Phosphate Problem of the Origin of Life. *Proc. Natl. Acad. Sci. USA* **2020**, *117* (2), 883–888.
- (93) Sasselov, D. D.; Grotzinger, J. P.; Sutherland, J. D. The Origin of Life as a Planetary Phenomenon. *Sci. Adv.* **2020**, *6* (6), eaax3419.
- (94) Yoneda, N.; Kusano, S.; Yasui, M.; Pujado, P.; Wilcher, S. Recent Advances in Processes and Catalysts for the Production of Acetic Acid. *Appl. Catal. A: General* **2001**, *221* (1), 253–265.
- (95) Berg, J. M.; Tymoczko, J. L.; Stryer, L. *Biochemistry*, 5th ed.; W. H. Freeman: New York, NY, 2002.
- (96) Weber, A. L. Origin of Fatty Acid Synthesis: Thermodynamics and Kinetics of Reaction Pathways. *J. Mol. Evol.* **1991**, *32* (2), 93–100.
- (97) Shock, E. L.; Schulte, M. D. Amino-Acid Synthesis in Carbonaceous Meteorites by Aqueous Alteration of Polycyclic Aromatic Hydrocarbons. *Nature* **1990**, *343* (6260), 728–731.
- (98) Sleep, N. H.; Zahnle, K. J.; Kasting, J. F.; Morowitz, H. J. Annihilation of Ecosystems by Large Asteroid Impacts on the Early Earth. *Nature* **1989**, *342* (6246), 139–142.
- (99) Marchi, S.; Bottke, W. F.; Elkins-Tanton, L. T.; Bierhaus, M.; Wuennemann, K.; Morbidelli, A.; Kring, D. A. Widespread Mixing and Burial of Earth's Hadean Crust by Asteroid Impacts. *Nature* **2014**, *511* (7511), 578–582.
- (100) Chyba, C.; Sagan, C. Endogenous Production, Exogenous Delivery and Impact-Shock Synthesis of Organic Molecules: An Inventory for the Origins of Life. *Nature* **1992**, *355* (6356), 125–132.
- (101) Bevan, A. W. R.; Bland, P. A.; Jull, A. J. T. Meteorite Flux on the Nullarbor Region, Australia. In *Meteorites: Flux with Time and Impact Effects*; Geological Society: London, Special Publications, 1998; Vol. 140, pp 59–73.

- (102) Macke, R. J.; Consolmagno, G. J.; Britt, D. T. Density, Porosity, and Magnetic Susceptibility of Carbonaceous Chondrites. *Meteoritics & Planetary Science* **2011**, *46* (12), 1842–1862.
- (103) Pierazzo, E.; Chyba, C. F. Impact Delivery of Prebiotic Organic Matter to Planetary Surfaces. In *Comets and the Origin and Evolution of Life*; Thomas, P. J., Hicks, R. D., Chyba, C. F., McKay, C. P., Eds.; Advances in Astrobiology and Biogeophysics; Springer: Berlin, Heidelberg, 2006; pp 137–168.
- (104) Keil, K. Mineralogical and Chemical Relationships among Enstatite Chondrites. *J. Geophys. Res.* **1968**, *73* (22), 6945–6976.
- (105) McDonough, W. F. Compositional Model for the Earth's Core. *Treatise on Geochemistry* **2003**, *2*, 547.
- (106) Dauphas, N. The Isotopic Nature of the Earth's Accreting Material through Time. *Nature* **2017**, *541* (7638), 521–524.
- (107) Bermingham, K. R.; Worsham, E. A.; Walker, R. J. New Insights into Mo and Ru Isotope Variation in the Nebula and Terrestrial Planet Accretionary Genetics. *Earth Planet Sci. Lett.* **2018**, *487*, 221–229.
- (108) Hellmann, J. L.; Hopp, T.; Burkhardt, C.; Becker, H.; Fischer-Gödde, M.; Kleine, T. Tellurium Isotope Cosmochemistry: Implications for Volatile Fractionation in Chondrite Parent Bodies and Origin of the Late Veneer. *Geochim. Cosmochim. Acta* **2021**, *309*, 313–328.
- (109) Macke, R. J.; Consolmagno, G. J.; Britt, D. T.; Hutson, M. L. Enstatite Chondrite Density, Magnetic Susceptibility, and Porosity. *Meteoritics & Planetary Science* **2010**, *45* (9), 1513–1526.
- (110) Lewis, J. S.; Prinn, R. G. *Planets and Their Atmospheres: Origin and Evolution*; International Geophysics Series; Academic Press Inc.: Orlando, FL, 1984; Vol. 33.
- (111) Wang, Z.; Mai, K.; Kumar, N.; Elder, T.; Groom, L. H.; Spivey, J. J. Effect of Steam During Fischer–Tropsch Synthesis Using Biomass-Derived Syngas. *Catal. Lett.* **2017**, *147* (1), 62–70.
- (112) Pendyala, V. R. R.; Shafer, W. D.; Jacobs, G.; Davis, B. H. Fischer–Tropsch Synthesis: Effect of Reaction Temperature for Aqueous-Phase Synthesis Over a Platinum Promoted Co/Alumina Catalyst. *Catal. Lett.* **2014**, *144* (6), 1088–1095.
- (113) Navarro-González, R.; Romero, A.; Honda, Y. Power Measurements of Spark Discharge Experiments. *Orig Life Evol Biosph* **1998**, *28* (2), 131–153.
- (114) Chiu, R.; Tinel, L.; Gonzalez, L.; Ciuraru, R.; Bernard, F.; George, C.; Volkamer, R. UV Photochemistry of Carboxylic Acids at the Air–Sea Boundary: A Relevant Source of Glyoxal and Other Oxygenated VOC in the Marine Atmosphere. *Geophys. Res. Lett.* **2017**, *44* (2), 1079–1087.
- (115) Simoneit, B. R. T.; Rushdi, A. I.; Deamer, D. W. Abiotic Formation of Acylglycerols under Simulated Hydrothermal Conditions and Self-Assembly Properties of Such Lipid Products. *Adv. Space Res.* **2007**, *40* (11), 1649–1656.
- (116) Rapf, R. J.; Perkins, R. J.; Dooley, M. R.; Kroll, J. A.; Carpenter, B. K.; Vaida, V. Environmental Processing of Lipids Driven by Aqueous Photochemistry of α -Keto Acids. *ACS Cent. Sci.* **2018**, *4* (5), 624–630.
- (117) Gibard, C.; Bhowmik, S.; Karki, M.; Kim, E.-K.; Krishnamurthy, R. Phosphorylation, Oligomerization and Self-Assembly in Water under Potential Prebiotic Conditions. *Nat. Chem.* **2018**, *10* (2), 212–217.
- (118) Sarkar, S.; Dagar, S.; Verma, A.; Rajamani, S. Compositional Heterogeneity Confers Selective Advantage to Model Protocellular Membranes during the Origins of Cellular Life. *Sci. Rep* **2020**, *10* (1), 4483.
- (119) Jia, T. Z.; Chandru, K.; Hongo, Y.; Afrin, R.; Usui, T.; Myojo, K.; Cleaves, H. J. Membraneless Polyester Microdroplets as Primordial Compartments at the Origins of Life. *Proc. Natl. Acad. Sci. USA* **2019**, *116* (32), 15830–15835.
- (120) Fares, H. M.; Marras, A. E.; Ting, J. M.; Tirrell, M. V.; Keating, C. D. Impact of Wet-Dry Cycling on the Phase Behavior and Compartmentalization Properties of Complex Coacervates. *Nat. Commun.* **2020**, *11* (1), 5423.

- (121) Dora Tang, T.-Y.; Rohaida Che Hak, C.; Thompson, A. J.; Kuimova, M. K.; Williams, D. S.; Perriman, A. W.; Mann, S. Fatty Acid Membrane Assembly on Coacervate Microdroplets as a Step towards a Hybrid Protocell Model. *Nat. Chem.* **2014**, *6* (6), 527–533.

Recommended by ACS

Meteorite Parent Body Aqueous Alteration Simulations of Interstellar Residue Analogs

Danna Qasim, Christopher K. Materese, *et al.*

JANUARY 09, 2023
ACS EARTH AND SPACE CHEMISTRY

READ 

Infrared Spectroscopy of (Benzene–H₂S–X_n)⁺, X = H₂O (n = 1 and 2) and CH₃OH (n = 1), Radical Cation Clusters: Microsolvation Effects on the S– π Hemibond

Takeru Kato and Asuka Fujii

JANUARY 13, 2023
THE JOURNAL OF PHYSICAL CHEMISTRY A

READ 

An Experimental and Theoretical Investigation of HCN Production in the Hadean Earth Atmosphere

Ben K. D. Pearce, Sarah M. Hörst, *et al.*

SEPTEMBER 23, 2022
ACS EARTH AND SPACE CHEMISTRY

READ 

Mineralochemical Mechanism for the Formation of Salt Volcanoes: The Case of Mount Dallol (Afar Triangle, Ethiopia)

Fermín Otálora, Juan Manuel García-Ruiz, *et al.*

OCTOBER 19, 2022
ACS EARTH AND SPACE CHEMISTRY

READ 

Get More Suggestions >